

LAEFF 95/13

FTUAM 95/..

RENORMALIZATION GROUP CALCULATION OF THE GALAXY-GALAXY CORRELATION FUNCTION

J. PÉREZ-MERCADER*, T. GOLDMAN[†], D. HOCHBERG^{‡,*a}
AND R. LAFLAMME[†]

**Laboratorio de Astrofísica Espacial y Física Fundamental
Apartado 50727, 28080 Madrid, Spain*

*[†]Theoretical Division, Los Alamos National Laboratory
Los Alamos, New Mexico 87545, USA*

*[‡]Departamento de Física Teórica, Universidad Autónoma de Madrid
Cantoblanco, 28049 Madrid, Spain*

The observable Universe is described by a collection of equal mass galaxies linked into a common unit by their mutual gravitational interaction. The partition function of this system is cast in terms of Ising model spin variables and maps exactly onto a three-dimensional stochastic scalar classical field theory. The full machinery of the renormalization group and critical phenomena is brought to bear on this field theory allowing one to calculate the galaxy-to-galaxy correlation function, whose critical exponent is predicted to be between 1.530 to 1.862, compared to the phenomenological value of 1.6 to 1.8

It is an empirical fact that the Universe today is dominated by matter which is in gravitational interaction and is organized into spatial configurations consisting of galaxies, groups of galaxies, superclusters, and even larger structures^{1,2}. It is also true that one of the outstanding problems in modern cosmology is to provide a quantitative description of this structure from first principles³. One of the standard ways in which structure, that is, the departure from randomness and homogeneity, is described, is by means of correlation functions. These objects are a basic tool for unravelling structure in physical systems ranging from gases and liquids to solids. Experimental input is typically deduced from analysis of the scattering of neutrons or x-rays. By contrast, in cosmology, this input is provided by direct observation. From observation, the galaxy-to-galaxy correlation function ($\xi_{gal}(r)$) is known to scale as $\xi(r) \sim r^{-\gamma}$, where $\gamma \sim O(1.6 \text{ to } 1.8)$, instead of $\gamma = 1$, which is what one would naively expect⁴ for a homogeneous distribution of matter in a three dimensional space.

^aBased on a talk presented by D. Hochberg at the International Workshop on Elementary Particle Physics, Valencia, Spain, June 5 to 9, 1995.

Given these observational facts, we shall demonstrate how to *calculate* the exponent γ from first principles⁵. We take our cue by analogy with the well established theory of liquids and interacting gases⁶, where the relative arrangements of the atoms and molecules (galaxies, in our case!) are accurately described in terms of the principles and methods of classical statistical mechanics. We will apply the same techniques to a system made up by many galaxies in gravitational interaction, subject to fluctuations emerging from the intrinsic properties of the gravitational interaction⁵. We will obtain the canonical partition function for this system and compute the two-point function and its corrections due to these fluctuations.

Begin by considering the continuous mass density $\rho(\mathbf{r})$ describing the spatial distribution of galaxies. The departure from perfect homogeneity is $\delta\rho(\mathbf{r}) = \rho(\mathbf{r}) - \bar{\rho}$, where $\bar{\rho}$ denotes the average density. The galaxy-galaxy correlation function is defined³ as $\xi_{Gal}(\mathbf{r}_i - \mathbf{r}_j) = \langle \delta\rho(\mathbf{r}_i)\delta\rho(\mathbf{r}_j) \rangle / \bar{\rho}^2$, the angular brackets denoting a suitable averaging. This is the joint probability distribution function for finding a galaxy at position $\mathbf{r}_i$ given that there is one at $\mathbf{r}_j$. In the static, nonrelativistic and weak gravitational field limit, (we discuss the effects of expansion later) the gravitational interaction energy for this system is

$$H_{int} = -\frac{G}{2} \int \int d^3\mathbf{r}_i d^3\mathbf{r}_j \rho(\mathbf{r}_i) \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|} \rho(\mathbf{r}_j), \quad (1)$$

where the integral excludes the point $\mathbf{r}_i = \mathbf{r}_j$, and the factor of 1/2 appears because each galaxy is counted twice in the integration. It is reasonable to consider the gas of N galaxies as made up of discrete spatially localized points of equal mass m_o at position $\mathbf{r}_j$, so that $\rho(\mathbf{r}) = \sum_{j=1}^N m_j \delta(\mathbf{r} - \mathbf{r}_j)$. The interaction energy then becomes

$$H_{int} = \frac{-1}{2} \sum_{i,j} \frac{m_i G m_j}{|\mathbf{r}_i - \mathbf{r}_j|} \equiv \frac{-1}{2} \sum_{i,j} m_i L_{ij}^{-1} m_j, \quad (2)$$

where m_j is the mass of the galaxy at site $\mathbf{r}_j$, and $i \neq j$. In this “granular” picture, the contrast function $\delta\rho(\mathbf{r}_i)/\bar{\rho}$ is equal to 1 if there is a galaxy at $\mathbf{r}_i$ and is equal to -1 if there is a void. Because we have assumed all N galaxies have equal mass, there is a natural correspondence between the m_i and a two-valued ($\pm$) “spin” variable s_i :

$$m_i = \frac{m_o}{2}(s_i + 1). \quad (3)$$

A galaxy at site $\mathbf{r}_i$ corresponds to “spin-up”, while a void or deficit corresponds to “spin-down”. Because of (3), we can express the Boltzmann factor and associated partition function of this gravitational Ising system in terms of the

spin variables as:

$$Z_{Ising}^{Grav}[\beta] = \sum_{\{s_i\}} \exp \left(\frac{1}{4} \sum_{i,j} s_i A_{ij}^{-1} s_j + \frac{1}{4} \sum_j s_j h_j \right), \quad (4)$$

where $A_{ij}^{-1} = \frac{\beta G m_o^2/2}{|\mathbf{r}_i - \mathbf{r}_j|}$, and $h_j = \sum_i \frac{\beta m_o^2 G}{|\mathbf{r}_i - \mathbf{r}_j|}$, and we have dropped an irrelevant constant term. β plays the role of an inverse temperature. The sum in (4) is over all possible spin configurations $s_1 = \pm 1, s_2 = \pm 1, \dots, s_N = \pm 1$.

To continue, we make use of the Hubbard-Stratonovich (or gaussian) transformation⁷

$$\begin{aligned} \pi^{N/2} (\det A)^{-1/2} \exp \left[\frac{1}{4} \sum_{i,j} s_i A_{ij}^{-1} s_j \right] = \\ \int_{-\infty}^{\infty} \Pi_{m=1}^N dx_m \exp \left(- \sum_{m,n} x_m A_{mn} x_n + \sum_m s_m x_m \right), \end{aligned} \quad (5)$$

which maps exactly the discrete system described by the spin variables s_i onto a system described by the continuous real variable $-\infty < x_m < \infty$. Applying (5) to the partition function in (4) yields

$$\begin{aligned} Z_{Ising}^{Grav}[\beta] &= C \int [d\phi] \exp \left[-\frac{\beta}{2} \int (\phi(\mathbf{r}) - h(\mathbf{r})) A(\mathbf{r}, \mathbf{r}') (\phi(\mathbf{r}') - h(\mathbf{r}')) d\mathbf{r} d\mathbf{r}' \right. \\ &\quad \left. + \int d\mathbf{r} \log \cosh(\beta \phi(\mathbf{r})) \right] \\ &\equiv \int [d\phi] e^{-\beta \mathcal{H}[\phi, h]}, \end{aligned} \quad (6)$$

performing the exact configurational sum over the $\{s_i\}$ and after taking the continuum limit which rigorously converts the discrete system into a continuous three dimensional 1-component scalar field theory. The linear term in (4) results in the highly nonlinear log cosh appearing in (6). C is an inessential factor and $h(\mathbf{r}) = \frac{1}{2} \int d\mathbf{r}' A^{-1}(\mathbf{r}, \mathbf{r}')$. The continuous scalar field ϕ is the order parameter for this system; its n -point Green functions correspond to the statistically averaged product of n times the contrast evaluated at n -sites. The operator A in the continuum limit is given by

$$A(\mathbf{r}, \mathbf{r}') = \delta^{(3)}(\mathbf{r} - \mathbf{r}') \frac{-\nabla^2}{2\pi\beta G m_o^2}, \quad (7)$$

since $\nabla^2 \frac{1}{|\mathbf{x}|} = -4\pi\delta^{(3)}(\mathbf{x})$.

As is well known⁸, the connected two-point function for the spin system and the field theory are the same. Moreover, the unavoidable fluctuations in the field ϕ result in an *anomalous* dimension η shifting the canonical dimension of the field such that when $|\mathbf{r} - \mathbf{r}'| \rightarrow \infty$, the two-point function for the hamiltonian in (6) scales as

$$\lim_{|\mathbf{r}-\mathbf{r}'| \rightarrow \infty} \langle s_i s_j \rangle = \lim_{|\mathbf{r}-\mathbf{r}'| \rightarrow \infty} \xi_{Gal}(|\mathbf{r} - \mathbf{r}'|) \sim |\mathbf{r} - \mathbf{r}'|^{-(d-2+\eta)}. \quad (8)$$

Here d is the dimension of space ($d = 3$) and η is the critical exponent for the pair correlation function whose value ($0.0198 - 0.064$) (see Table 1) differs from zero due to the fluctuations in ϕ .

Thus, our calculation shows that for large separations, the galaxy-galaxy correlation function must scale as

$$\xi_{Gal}(|\mathbf{r}_i - \mathbf{r}_j|) \sim r^{-(d-2+\eta)}, \quad (9)$$

with⁹ $1.0198 \leq 1 + \eta \leq 1.064$ for $d = 3$, and without taking the background expansion into account.

Thus far, our calculation has been performed for a static ensemble of galaxies, but the Universe is expanding and the effects of this expansion will modify the values of the critical exponents. From the theory of dynamical critical phenomena, it is phenomenologically known from computer simulations that the way time enters into the correlation function is by a modification of its argument, viz.,

$$\xi(r) \rightarrow \xi(r; t) = F(r/l(t)), \quad (10)$$

where $l(t) \sim t^\zeta$ is a function of time and here, r denotes the comoving coordinate, $r_{comoving}$, which is related to the physical coordinate via

$$r_{physical} = a(t) r_{comoving}, \quad (11)$$

with $a(t)$ the scale factor for the background spacetime. From computer simulations in condensed matter physics, it is established that $\zeta = 1/3$ for systems with a conserved order parameter, while $\zeta = 1/2$ when the order parameter is not conserved.

For a matter dominated expansion, the case we are interested in, $a(t) = t^{2/3}$ and thus for a given physical separation

$$t \sim (r_{comoving})^{-3/2}. \quad (12)$$

This means that the function $l(t)$ scales as

$$l(t) = t^\zeta = (r_{comoving})^{-3\zeta/2}. \quad (13)$$

Table 1: Calculated values for the galaxy-galaxy correlation function critical exponent

Method of Calculation	γ_{Static}	$\gamma_{Expanding}^C$	$\gamma_{Expanding}^{NC}$
Series estimates	1.056 ± 0.008	1.584 ± 0.012	1.848 ± 0.014
$O(\epsilon)$	0	1.5	1.75
$O(\epsilon^2)$	1.0198	1.530	1.785
$O(\epsilon^3)$	1.037	1.555	1.815
$O(\epsilon^4)$	1.029	1.543	1.801

Putting together these facts yields the scaling behavior of $\xi(r; t)$:

$$\begin{aligned}
 \xi(r; t) &= \left(r_{comoving} / r_{comoving}^{-3\zeta/2} \right)^{-(d-2+\eta)} \\
 &= (r_{comoving})^{-(d-2+\eta)(1+3\zeta/2)}.
 \end{aligned} \tag{14}$$

In other words, the critical exponent resulting from the expansion of the Universe and dynamical critical phenomena effects is

$$\gamma = (d - 2 + \eta) \times (1 + 3\zeta/2). \tag{15}$$

The predicted (calculated) value for γ is between 1.530 and 1.596 for $\zeta = 1/3$ (for a conserved order parameter) and between 1.75 to 1.862 for $\zeta = 1/2$ (non-conserved order parameter). Table 1 displays these exponents for both the static and expanding cases (C, NC, denote conserved and nonconserved order parameter, respectively). The static values are taken from Ref[9]. The last four entries refer to expansions in powers of $\epsilon = 4 - d$. These values are to be compared with the values inferred from current galaxy catalogs, which range from 1.5 for the APM survey, to 1.8 for the Lick survey^{2,1}.

We have thus accomplished what we set out to do, namely, provide a first-principles calculation of the galaxy-galaxy correlation function using standard methods of statistical mechanics and the theory of critical phenomena. Naturally, one might (and should) ask *why* this calculation is successful, in other words, what does a system of galaxies have to do with a spin model? The answer is provided by the universality hypothesis, according to which rather distinct physical systems will exhibit *identical* critical behavior provided the systems in question possess the same space dimensionality and have order parameters with the same dimensionality (or number of independent components). We see that this is the case here: $d = 3$ for both systems, while the scalar field and the magnetization are one-component order parameters for the galaxy system and the 3d Ising model, respectively. In other words, both

systems belong to the same universality class. Thus, we must expect them to have the same values for their critical exponents.

Acknowledgements

D.H. would like to thank the organizers for the invitation to participate in this Workshop and for their kind and generous hospitality. Thanks are also due to the participants for their enthusiastic response to this talk, as evidenced by their many thoughtful comments and questions.

References

1. The Lick survey is based on plates taken by C.D. Shane and C.A. Wirtanen, *Publ. Lick Obs.* **22** Part 1 (1967).
2. S.J. Maddox et al., MNRAS **242** (1990) 43P.
3. See, e.g., P.J.E. Peebles, *Principles of Physical Cosmology* (Princeton University Press, Princeton, 1993)
4. As we shall demonstrate below, the galaxy-galaxy correlation function can be expressed as a scalar field propagator. In units of length, a scalar field in d -dimensional space has a canonical dimension of $-(d/2 - 1)$. The above statement about γ follows at once.
5. J. Pérez-Mercader, T. Goldman, D. Hochberg and R. Laflamme, Los Alamos preprint LA-UR-95-1055.
6. See, e.g., D.L. Goodstein, *States of Matter* (Dover, New York, 1985).
7. See, e.g., J. Binney, N. Dowrick, A. Fisher and M. Newman, *The Theory of Critical Phenomena: An Introduction to the Renormalization Group*, (Oxford University Press, Oxford, 1992). Also, D. Amit, *Field Theory, the Renormalization Group and Critical Phenomena*, revised second edition, (World Scientific, Singapore, 1984).
8. K.G. Wilson, Phys. Rev. **179**, 1499 (1969).
9. M.N. Barber, Phys. Rep. **29C**, 1 (1977).